Nine-Nina simulations and predictions over 1983-1993 f r o m Cane and Zebiak's m o d e l and satellite observations.

- B. Dewitte, <u>C Perigaud</u> and Y. Chao (Jet Propulsion laboratory, California institute of Technology, 4800 oak Grove Drive, Pasadena, CA 91109, USA; cp@pacific.jpl.nasa.gov)
- Anomalies of thermocline depths, SS"1', winds and cloud convection relative to monthly climatology are first simulated with the Cane and Zebiak's model in its uncoupled mode forced by FSU wind anomalies. Simulations are compared with Geosat sea-level, NOAA/AVIIRRSST, FSU winds and ISCCP cloud convection observations. Mode] and data are in good agreement for sea-level, but mode] fails to simulate SS"1', wind and cloud Nina anomaly in 1988.
- SS'1', cloud convection and air-sea heat budgets derived from SSM1 are then used to force the atmospheric part and/or mixed layer part of the model in its uncoupled mode. Simulations are then considerably improved. But used in its coupled mode, discrepancy shows up again because of the importance of the initial conditions in predictions.
- Twin experiments are then runto study the impact of initializing predictions with sea-level or with SS'1'. Impact of sca-level on predictions lasts longer than that of SST. Predictions are degraded or improved, depending on the way the coupled model is further corrected or not with extra observations during the predictions.

- 1. 1994 Ocean Sciences meeting
- 2. 008617393
- 3. (a) C. Perigaud
 Jet Propulsion Laboratory,
 MS 300/323
 Pasadena, CA 91109
 (b) tel:818- 354-8203
 (C) fax: 818-393-6720
- 4, **U**
- 5.
- (a) 4504: air-sea interaction
- (b) 4231: equatorial Oceanography
- (c) 4215: Climate and interannual variability
- 6.0
- 7. 0?40
- 8. Billing to:
 Claire M PERIGAUD
 VISA n": 4070"009002000170
 expires 06/94
- 9. C
- 1 (). none 11. No.